

Spatio-temporal dynamic modeling of plant communities responses to hydrological pressures in a semiarid Mediterranean wetland

J. Martínez-López¹, J. Martínez-Fernández^{1,2}, B. Naimi³,
M.F. Carreño¹ and M.A. Esteve¹

¹Ecology and Hydrology Department - University of Murcia (Murcia, Spain)

²Applied Biology Dept. University Miguel Hernandez (Elche, Spain)

³ITC - University of Twente (Enschede, The Netherlands)

October 28th - 31st



isem 2013
TOULOUSE - FRANCE

Study area


Wetland model

ISEM 2013

Introduction

Methods

Conclusions



Marina del Carmoli wetland (300 ha)

Wetland model

ISEM 2013

Introduction

Methods

Conclusions



Wetland plant communities

Semiarid Mediterranean saline wetlands are semi-terrestrial ecosystems

HUMIDITY




SALINITY

- ▶ Salt steppe (left) - priority habitat by the Habitats Directive
- ▶ Salt marsh (center) - habitat of interest by the HD
- ▶ Reed beds (right) (*Phragmites australis*) - invasive


External water inputs

Percentage of irrigated areas has increased in the last decades due to the opening of a water transfer (Martínez-López et al., 2013)



Plant communities change

Important plant communities are being lost!




Carreño et al., 2008; Martínez-López et al., 2012

- ▶ Spatially explicit wetland model of how irrigated agriculture is affecting plant community composition in this semiarid Mediterranean wetland

- ▶ R as a modelling environment:
 - ▶ GIS capabilities
 - ▶ source code is flexible
 - ▶ free availability and growing user community




- ▶ Wetland is divided into pixels (25 m)
- ▶ Plant communities are modelled separately pixel by pixel (4 maps)
- ▶ The total abundance of plant communities within a pixel is limited so:
 - ▶ competition among plant communities mediated by
 - ▶ total drainage water input to the wetland
 - ▶ spatial environmental variables influencing water availability and growth
 - ▶ the dispersion of other PC from the surrounding pixels



Initial and validation maps of plant communities

Model was tested by means of remote sensing data for the period 1992-2008



Carreño et al., 2008; Martínez-López et al., 2012


ISEM 2013



Model assumptions II

native vs. invasive taxa

- ▶ invasive reed beds are potentially present in all pixels
- ▶ salt marsh is able to disperse into neighbour pixels



Non spatial forcing input

Wetland model

ISEM 2013


Drainage water input

WARP index (Martínez-López et al., 2014a,b)

Introduction

Methods

Conclusions



Wetland environmental spatial parameters

Wetland model


ISEM 2013

Introduction

Methods


Conclusions

- ▶ (A) distance map to ephemeral river 1 (**reed beds**)
- ▶ (B) distance map to ephemeral river 2 (**reed beds**)
- ▶ (C) Flow accumulation map (**salt marsh**)



- ▶ All parameters are on a relative 0–1 scale.

Model diagram



1. Initial dynamic model was developed using Stella (1 pixel)
2. Conversion to R using 'StellaR' script (Naimi and Voinov, 2012)
3. State variables and spatial environmental variables as matrices
4. Model wrapped as a R function
5. `ode.2D("euler" method, time = 24 year, TS = 0.25)` (library "deSolve")

1. The model serves as a tool for
 - ▶ wetland conservation and management studies (habitat loss)
 - ▶ testing plant community interactions
 - ▶ testing relationships between plant communities and environmental variables in space and time
2. The library undergoes further developments in order to become a flexible tool for the development of new spatio-dynamic models