

Package ‘MWright’

January 20, 2025

Type Package

Title Mainardi-Wright Family of Distributions

Version 0.3.2

Author Dexter Cahoy

Maintainer Dexter Cahoy <cahoyd@uhd.edu>

Description Implements random number generation, plotting, and estimation algorithms for the two-parameter one-sided and two-sided M-Wright (Mainardi-Wright) family.

The M-Wright distributions naturally generalize the widely used one-sided (Airy and half-normal or half-Gaussian) and symmetric (Airy and Gaussian or normal) models.

These are widely studied in time-fractional differential equations. References: Cahoy and Minkabu (2017) <[doi:10.3233/MAS-170388](https://doi.org/10.3233/MAS-170388)>; Cahoy (2012) <[doi:10.1007/s00180-011-0269-x](https://doi.org/10.1007/s00180-011-0269-x)>; Cahoy (2012) <[doi:10.1080/03610926.2010.543299](https://doi.org/10.1080/03610926.2010.543299)>; Cahoy (2011); Mainardi, Mura, and Pagnini (2010) <[doi:10.1155/2010/104505](https://doi.org/10.1155/2010/104505)>.

License GPL (>= 3)

Encoding UTF-8

LazyData true

Imports stats, cubature

RoxxygenNote 6.1.1

NeedsCompilation no

Repository CRAN

Date/Publication 2019-08-07 23:00:05 UTC

Contents

dmwright1	2
dmwright2	3
int_est1	4
int_est2	5
MWright	6
pmwright1	6
pmwright2	7

point_est1	8
point_est2	9
rmwright1	10
rmwright2	11

Index	13
--------------	-----------

dmwright1	<i>One-sided M-Wright distribution</i>
-----------	--

Description

Plots the density function.

Usage

```
dmwright1(ah, sh, m, max)
```

Arguments

ah	point estimate for shape parameter alpha.
sh	point estimate for scale parameter s.
m	number of data points (pairs) to use for plotting.
max	maximum x-axis value to use for plotting.

Value

numeric matrix

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
xy=dmwright1(0.45, 2.5, 1000, 10)
plot(xy[,1], xy[,2], lwd = 2, type="l", ylab="", xlab="x")

mwright1_sided <- rmwright1(1000, 0.45, 2.5)
hist(mwright1_sided, br=30, prob=TRUE)
lines(xy[,1], xy[,2], lwd=2 )
```

dmwright2

Two-sided M-Wright distribution

Description

Plots the density function.

Usage

```
dmwright2(ah, sh, m, max)
```

Arguments

ah	point estimate for shape parameter alpha.
sh	point estimate for scale parameter s.
m	number of data points (pairs) to use for plotting.
max	maximum x-axis value to use for plotting.

Value

numeric matrix

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics- Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
xy=dmwright2(0.45, 2.5, 1000, 10)
plot(xy[,1], xy[,2], lwd = 2, type="l",ylab="", xlab="x")

mwright2_sided <- rmwright2(1000, 0.45, 2.5)
hist(mwright2_sided, br=30, prob=TRUE)
lines(xy[,1], xy[,2], lwd=2 )
```

int_est1

Interval estimation for one-sided M-Wright distribution

Description

Confidence intervals for the model parameters.

Usage

```
int_est1(x, lev)
```

Arguments

x	numeric vector
lev	confidence level.

Value

matrix

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics- Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
mwright_1sided <- rmwright1(1000, 0.7, 0.4)
int_est1(mwright_1sided ,0.95)
```

int_est2

Interval estimation for two-sided M-Wright distribution

Description

Confidence intervals for the model parameters.

Usage

```
int_est2(x, lev)
```

Arguments

x	numeric vector
lev	confidence level.

Value

matrix

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics-Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
mwright_2sided <- rmwright2(1000, 0.7, 0.4)
int_est2(mwright_2sided ,0.95)
```

MWright*MWright Package***Description**

Contains random number generation, plotting, and estimation algorithms for the two-parameter one-sided and two-sided M-Wright (Mainardi-Wright) family. The M-Wright distributions naturally generalize widely used one-sided (Airy and half-normal or half-normal) and symmetric (Airy and Gaussian or normal) models. These are widely studied in time-fractional diffusion processes.

Details

References:

- Cahoy and Minkabo (2017)<doi:10.3233/MAS-170388>
- Cahoy (2012) <doi:10.1007/s00180-011-0269-x>
- Cahoy (2012) <doi:10.1080/03610926.2010.543299>
- Cahoy (2011)
- Mainardi, Mura, and Pagnini (2010) <doi:10.1155/2010/104505>

Author(s)

Dexter Cahoy <cahoyd@uhd.edu>

pmwright1*Distribution function for one-sided M-Wright distribution***Description**

Calculates a left-tail probability.

Usage

```
pmwright1(alp, sc, upper)
```

Arguments

- | | |
|--------------|---|
| alp | point estimate for shape parameter alpha. |
| sc | point estimate for scale parameter s. |
| upper | non-negative upper quantile |

Value

numeric

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics- Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
pmwright1(runif(1), runif(1,0,10),Inf )
pmwright1(runif(1), runif(1,0,10), 0.5 )
```

pmwright2

Distribution function for two-sided M-Wright distribution

Description

Calculates a left-tail probability.

Usage

```
pmwright2(alp, sc, upper)
```

Arguments

alp	point estimate for shape parameter alpha.
sc	point estimate for scale parameter s.
upper	upper quantile

Value

numeric

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics-Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
pmwright2(runif(1), runif(1,0,10), Inf )
pmwright2(runif(1), runif(1,0,10), 0.5 )
```

point_est1

Point estimation for one-sided M-Wright distribution

Description

This provides point estimates for the shape and scale parameters.

Usage

```
point_est1(x)
```

Arguments

x	numeric vector.
---	-----------------

Value

numeric vector

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics- Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
x <- rmwright1(1000, 0.7, 0.4)
point_est1(x)
```

point_est2

Point estimates for two-sided M-Wright distribution

Description

This provides point estimates for the shape and scale parameters.

Usage

```
point_est2(x)
```

Arguments

x numeric vector.

Value

numeric vector

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics-Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
x <- rmwright2(1000, 0.7, 0.4)
point_est2(x)
```

rmwright1

Random number generation for one-sided M-Wright distribution

Description

Generates random numbers.

Usage

```
rmwright1(n, nu, sc)
```

Arguments

- n sample size.
- nu a number between 0 and 1.
- sc a non-negative scale value.

Value

a vector of one-sided M-Wright distributed random numbers

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics-Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoy (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
mrwright_1sided <- rmwright1(1000, 0.7, 0.4)
hist(mwright_1sided, br=30)
```

rmwright2

Random number generation for two-sided M-Wright distribution

Description

Generates random numbers.

Usage

```
rmwright2(n, nu, sc)
```

Arguments

- n sample size.
- nu a number between 0 and 1.
- sc a non-negative scale value.

Value

a vector of two-sided M-Wright distributed random numbers.

References

- Cahoy and Minkabo (2017). *Inference for three-parameter M-Wright distributions with applications*. Model Assisted Statistics and Applications, 12(2), 115-125. <https://doi.org/10.3233/MAS-170388>
- Cahoy (2012). *Moment estimators for the two-parameter M-Wright distribution*. Computational Statistics, 27(3), 487-497. <https://doi.org/10.1007/s00180-011-0269-x>
- Cahoy (2012). *Estimation and simulation for the M-Wright function*. Communications in Statistics- Theory and Methods, 41(8), 1466-1477. <https://doi.org/10.1080/03610926.2010.543299>
- Cahoyd (2011). *On the parameterization of the M-Wright function*. Far East Journal of Theoretical Statistics, 34(2), 155-164. <http://www.pphmj.com/abstract/5767.htm>
- Mainardi, Mura, and Pagnini (2010). *The M-Wright Function in Time-Fractional Diffusion Processes: A Tutorial Survey*. Int. J. Differ. Equ., Volume 2010. <https://doi.org/10.1155/2010/104505>

Examples

```
mwright_2sided <- rmwright2(1000, 0.7, 0.4)
hist(mwright_2sided, br=30)
```

Index

`dmwright1`, 2
`dmwright2`, 3

`int_est1`, 4
`int_est2`, 5

`MWright`, 6
`MWright-package (MWright)`, 6

`pmmwright1`, 6
`pmmwright2`, 7
`point_est1`, 8
`point_est2`, 9

`rmwright1`, 10
`rmwright2`, 11